

The equality problem for infinite words generated by primitive morphisms

Juha Honkala
 Department of Mathematics
 University of Turku
 20014 Turku, Finland
 email: juha.honkala@utu.fi

Abstract

We study the equality problem for infinite words obtained by iterating morphisms. In particular, we give a practical algorithm to decide whether or not two words generated by primitive morphisms are equal.

1 Introduction

The equality problem for pure morphic words (also called the D0L ω -equivalence problem) was solved by Culik II and Harju [2]. A simpler approach was suggested in [6]. The algorithms given in [2, 6] have very high complexities. No practical algorithm is known in the general case. For the equality problem of infinite words generated by polynomially bounded morphisms see [3].

The purpose of this paper is to give a practical algorithm for the equality problem of pure morphic words generated by primitive morphisms. By definition, a morphism $g : X^* \rightarrow X^*$ is primitive if there is a positive integer k such that for all $a, b \in X$, a occurs in $g^k(b)$.

To explain our result let X be a finite alphabet and let $g : X^* \rightarrow X^*$ be a morphism. Let $x \in X$ be a letter such that x is a prefix of $g(x)$. Then $g^i(x)$ is a prefix of $g^{i+1}(x)$ for all $i \geq 0$. If the length of $g^{i+1}(x)$ is always greater than the length of $g^i(x)$, we let $g^\omega(x)$ be the infinite word which has prefix $g^i(x)$ for all $i \geq 0$. Infinite words generated in this way by iterating a morphism are called pure morphic words (or infinite D0L words).

Let now X be an alphabet having $n \geq 2$ letters. Define $A(n) = \lfloor 9n^3 \sqrt{n \log n} \rfloor$. Let $g : X^* \rightarrow X^*$ and $h : X^* \rightarrow X^*$ be primitive morphisms and let $x \in X$ be a letter such that $g^\omega(x)$ and $h^\omega(x)$ exist. Define $f_1 = g^{2n-2}h^{2n-2}$ and $f_2 = h^{2n-2}g^{2n-2}$. We will show that $g^\omega(x) = h^\omega(x)$ if and only if f_1 and f_2 satisfy a balance condition and one of the words $f_1 f_1^{A(n)}(x)$ and $f_2 f_1^{A(n)}(x)$ is a prefix of the other.

The paper is structured as follows. In Section 2 we recall some basic definitions and earlier results. In particular, we define looping and loop-free mor-

phisms. Intuitively, a morphism $g : X^* \longrightarrow X^*$ is looping if some power of g generates a periodic word. In Section 3 we recall the reduction of the equality problem to the comparability problem. In Section 4 we study balance properties of morphisms. In Section 5 we solve the equality problem for loop-free primitive morphisms by using ideas and results from [5]. In Section 6 we solve the equality problem for looping primitive morphisms. The main result is presented in Section 7.

We assume that the reader is familiar with the basics concerning free monoid morphisms and their iterations, see [1, 9, 10]. For morphic words and their applications see also [8].

2 Preliminaries

2.1 Basic definitions

We use standard language-theoretic notation and terminology. In particular, the *length* of a word w is denoted by $|w|$. If $w \in X^*$ is a nonempty word, then $\text{alph}(w)$ is the set of all letters of X occurring in w . If w is a nonempty word, then w^ω is the infinite word

$$w^\omega = www \dots$$

If $u, v \in X^*$ are words, v is a *factor* of u if there exist words $u_1, u_2 \in X^*$ such that $u = u_1vu_2$. If, furthermore, $u_1 = \varepsilon$ then v is a *prefix* of u . If q is a nonnegative integer and w is a finite word or an infinite word, then $\text{Pref}_q(w)$ is the prefix of length q of w and $F_q(w)$ is the set of factors of length q of w . If $|w| < q$, it is understood that $\text{Pref}_q(w) = w$. If $L \subseteq X^*$, then

$$\text{Pref}_q(L) = \{\text{Pref}_q(w) \mid w \in L\}$$

and

$$F_q(L) = \bigcup_{w \in L} F_q(w).$$

Let $u \in X^*$ be a nonempty word and denote $u = a_1 \dots a_k$ where $a_i \in X$ for $i = 1, \dots, k$. Then a positive integer p is called a *period* of u if

$$a_{p+i} = a_i \quad \text{for } i = 1, \dots, k - p. \quad (1)$$

The smallest positive integer p satisfying (1) is called *the period* of u and is denoted by $\text{PER}(u)$.

Let $g : X^* \longrightarrow X^*$ be a morphism and let $w \in X^*$ be a word. If w is a prefix of $g(w)$, then $g^n(w)$ is a prefix of $g^{n+1}(w)$ for all $n \geq 0$. If, furthermore,

$$\lim_{n \rightarrow \infty} |g^n(w)| = \infty,$$

then we denote

$$g^\omega(w) = \lim_{n \rightarrow \infty} g^n(w).$$

In all other cases $g^\omega(w)$ is not defined. Hence, if $g^\omega(w)$ exists, it is the unique infinite word which has the prefix $g^n(w)$ for all $n \geq 0$.

2.2 Primitive morphisms

Suppose $g : X^* \longrightarrow X^*$ is a morphism. Then g is called *primitive* if there exists a positive integer k such that for all $x, y \in X$, y occurs in $g^k(x)$. If the alphabet X contains at least two letters, primitive morphisms are growing morphisms. In general, a morphism $g : X^* \longrightarrow X^*$ is called *growing* if for every $x \in X$ we have

$$|g^i(x)| \rightarrow \infty \quad \text{when} \quad i \rightarrow \infty.$$

If $g : X^* \longrightarrow X^*$ is a morphism, define

$$M_g = \max\{|g(x)| \mid x \in X\}$$

and

$$\text{CYCLIC}(g) = \{x \in X \mid g(x) \text{ contains at least one occurrence of } x\}.$$

The following lemma gives some basic properties of primitive morphisms.

Lemma 1 *Let X be an alphabet having $n \geq 2$ letters and let $g : X^* \longrightarrow X^*$ be a primitive morphism. Then*

- (i) $|g^n(x)| \geq M_g$ for all $x \in X$.
- (ii) If $\text{CYCLIC}(g) \neq \emptyset$, then $\text{alph}(g^{2n-2}(x)) = X$ for all $x \in X$.

Proof. Let $x, y \in X$. Because g is primitive, there is an integer i such that y occurs in $g^i(x)$ and $i \leq n - 1$. Hence $|g^n(x)| \geq |g(y)|$. This implies (i).

Assume then that $z \in \text{CYCLIC}(g)$. Then $\text{alph}(g^{n-1}(z)) = X$. Furthermore, if $x \in X$, then z occurs in $g^{n-1}(x)$. These facts imply (ii). $\square$

For the proof of the following lemma see [4].

Lemma 2 *Let X be an alphabet with $n \geq 2$ letters and let $g : X^* \longrightarrow X^*$ be a growing morphism. Let $w \in X^*$. Then*

$$\text{Pref}_2(\{g^i(w) \mid i \geq 0\}) = \text{Pref}_2(\{g^i(w) \mid i = 0, 1, \dots, 3n - 2\})$$

and

$$F_3(\{g^i(w) \mid i \geq 0\}) = F_3(\{g^i(w) \mid i = 0, 1, \dots, 2n^2 + 2n - 3\}).$$

2.3 Looping and loop-free morphisms

Let $g : X^* \longrightarrow X^*$ be a morphism. Then we say that g is *looping* if there exist a positive integer k , a letter $x \in X$ and a nonempty word u such that $(g^k)^\omega(x)$ exists and

$$(g^k)^\omega(x) = u^\omega.$$

If g is not looping, we say that g is *loop-free*.

Loop-free morphisms avoid small periods in the sense of the following lemma [5].

Lemma 3 *Let X be an alphabet having n letters and let $g : X^* \longrightarrow X^*$ be a morphism. If g is growing and loop-free, then*

$$\text{PER}(g^d(x)) \geq \frac{1}{2M_g^n} |g^d(x)|$$

whenever $x \in X$ and d is a positive integer.

For the proof of the next lemma see [7].

Lemma 4 *Let X be an alphabet having n letters and let $g : X^* \longrightarrow X^*$ be a primitive morphism. Assume that $x \in X$, k is a positive integer and*

$$(g^k)^\omega(x) = u^\omega,$$

where $u \in X^$ is a primitive word. Then*

$$|g^{2n}(z)| \geq 2|u|$$

for all $z \in X$.

3 Properties of infinite words

In this section we recall some results concerning infinite words generated by morphisms.

Let $g : X^* \longrightarrow X^*$ be a morphism and let $x \in X$ be a letter such that $g^\omega(x)$ exists. If w is a nonempty prefix of $g^\omega(x)$, then w is a proper prefix of $g(w)$.

Lemma 5 *Let $g_i : X^* \longrightarrow X^*$, $i = 1, 2$, be morphisms and let $x \in X$ be a letter such that $g_i^\omega(x)$ exists for $i = 1, 2$. Suppose*

$$g_i^\omega(x) = wb_i \dots$$

for $i = 1, 2$, where $w \in X^$, $b_i \in X$ and $b_1 \neq b_2$. If $i_1, \dots, i_k \in \{1, 2\}$, then either $g_{i_k} \dots g_{i_1}(x)$ is a prefix of w or has prefix wb_{i_k} .*

Proof. Consider the word $g_{i_k} \dots g_{i_1}(x)$ and assume that the claim holds for $g_{i_{k-1}} \dots g_{i_1}(x)$. If $g_{i_{k-1}} \dots g_{i_1}(x)$ is a prefix of w , the claim holds. Assume w is prefix of $g_{i_{k-1}} \dots g_{i_1}(x)$. Then $g_{i_k}(w)$ is a prefix of $g_{i_k} \dots g_{i_1}(x)$. The claim follows because $g_{i_k}(w)$ is a prefix of $g_{i_k}^\omega(x)$ which is longer than w . $\square$

Lemma 6 *Let $g : X^* \longrightarrow X^*$ and $h : X^* \longrightarrow X^*$ be morphisms and let $x \in X$ be a letter such that $g^\omega(x)$, $h^\omega(x)$, $(gh)^\omega(x)$ and $(hg)^\omega(x)$ exist. Then*

$$g^\omega(x) = h^\omega(x) \tag{2}$$

if and only if

$$(gh)^\omega(x) = (hg)^\omega(x). \tag{3}$$

Proof. Assume first that (2) holds. If $u \in \text{Pref}(g^\omega(x))$, then $gh(u) \in \text{Pref}(g^\omega(x))$. Hence

$$(gh)^i(x) \in \text{Pref}(g^\omega(x))$$

for all $i \geq 0$. Therefore

$$(gh)^\omega(x) = g^\omega(x).$$

Similarly

$$(hg)^\omega(x) = g^\omega(x).$$

Hence (3) holds.

Assume then that (2) does not hold. Let

$$g^\omega(x) = wa \cdots, \quad h^\omega(x) = wb \cdots,$$

where $w \in X^*$, $a, b \in X$ and $a \neq b$. Then for large values of i by Lemma 5, wa is a prefix of $(gh)^i(x)$ and wb is a prefix of $(hg)^i(x)$. Hence (3) does not hold. $\square$

Let $g : X^* \rightarrow X^*$ and $h : X^* \rightarrow X^*$ be morphisms. Then the set $\text{COMP}(g, h)$ is defined by

$$\begin{aligned} \text{COMP}(g, h) &= \{w \in X^* \mid \text{there exist } u_1, u_2 \in X^* \text{ such that} \\ &\quad g(w)u_1 = h(w)u_2\}. \end{aligned}$$

Lemma 7 *Let $g : X^* \rightarrow X^*$ and $h : X^* \rightarrow X^*$ be morphisms and let $x \in X$ be a letter such that $g^\omega(x)$ and $h^\omega(x)$ exist. Then*

$$g^\omega(x) = h^\omega(x) \tag{4}$$

if and only if

$$\text{Pref}(g^\omega(x)) \subseteq \text{COMP}(g, h). \tag{5}$$

Proof. Assume first that (4) holds. Assume that w is a prefix of $g^\omega(x) = h^\omega(x)$. Then $g(w)$ is a prefix of $g^\omega(x)$ and $h(w)$ is a prefix of $h^\omega(x)$. Hence $w \in \text{COMP}(g, h)$.

Assume then that (4) does not hold. Let

$$g^\omega(x) = wa \cdots, \quad h^\omega(x) = wb \cdots,$$

where $w \in X^*$, $a, b \in X$ and $a \neq b$. Then $g(w)$ has prefix wa and $h(w)$ has prefix wb . Because $a \neq b$, the word w is not in $\text{COMP}(g, h)$ which shows that (5) does not hold. $\square$

4 Balance properties of morphisms

Let $g : X^* \rightarrow X^*$ and $h : X^* \rightarrow X^*$ be morphisms. Define

$$\text{BAL}(g, h) = \max\{|gg^k(x)| - |hg^k(x)| \mid x \in X, k \geq 0\},$$

where the right-hand side is a nonnegative integer or ∞ .

For the proof of the following lemma see [4].

Lemma 8 *Let X be an alphabet having $n \geq 2$ letters and let $g : X^* \rightarrow X^*$ and $h : X^* \rightarrow X^*$ be morphisms. Define $M = \max\{M_g, M_h\}$. If*

$$\text{BAL}(g, h) < \infty,$$

then

$$\text{BAL}(g, h) \leq M^{2n-1} \exp(n^2(1 + \sqrt{6n \log n})).$$

Next we recall a very important result due to Culik II and Harju [2]. Let $g : X^* \rightarrow X^*$ be a morphism and let $a \in X$ be a letter. Define $X_1 = X - \{a\}$. Then the triple (X, g, a) is called a *1-system* if

- (i) $g(a) \in aX_1^*$,
- (ii) $g(x) \in X_1^*$ whenever $x \in X_1$,
- (iii) $\{g^i(a) \mid i \geq 0\}$ is infinite,
- (iv) if $x \in X_1$, then x occurs infinitely many times in $g^\omega(a)$.

A 1-system (X, g, a) is called *1-simple* if the restriction of g to X_1 is primitive.

For the proof of the following result see [2].

Theorem 9 *Let (X, g_i, a) , $i = 1, 2$, be 1-systems such that $g_1^\omega(a) = g_2^\omega(a)$. If (X, g_1g_2, a) and (X, g_2g_1, a) are 1-simple then there exists a positive integer K such that*

$$||g_1g_2(w)| - |g_2g_1(w)|| \leq K$$

whenever w is a prefix of $g_1^\omega(a)$.

In what follows we need the following consequence of Theorem 9.

Lemma 10 *Let X_1 be a finite alphabet. Let $g_1 : X_1^* \rightarrow X_1^*$ and $g_2 : X_1^* \rightarrow X_1^*$ be primitive morphisms and let $x \in X_1$ be a letter such that $g_1^\omega(x)$ and $g_2^\omega(x)$ exist and are equal. If g_1g_2 and g_2g_1 are primitive then there exists a positive integer K such that*

$$||g_1g_2(w)| - |g_2g_1(w)|| \leq K$$

whenever w is a factor of $g_1^\omega(x)$.

Proof. It is enough to prove the claim for the prefixes of $g_1^\omega(x)$. This claim is a consequence of Theorem 9. Indeed, choose a new letter $a \notin X_1$ and define $X = X_1 \cup \{a\}$. Assume that $g_i(x) = xu_i$ ($i = 1, 2$) and extend g_i by $g_i(a) = au_i$ ($i = 1, 2$). Then (X, g_i, a) , $i = 1, 2$, are 1-systems and $g_1^\omega(a) = g_2^\omega(a)$. Furthermore, (X, g_1g_2, a) and (X, g_2g_1, a) are 1-simple. Hence Theorem 9 is applicable. $\square$

5 The equality problem for loop-free primitive morphisms

5.1 The comparability problem

In this subsection we first recall a result from [5].

Let X be an alphabet having $n \geq 2$ letters and let $g : X^* \longrightarrow X^*$ and $h : X^* \longrightarrow X^*$ be growing morphisms. Define the mapping $\beta : X^* \longrightarrow \mathbb{Z}$ by

$$\beta(w) = |g(w)| - |h(w)|, \quad w \in X^*.$$

Assume that $t \geq 2$ is a fixed integer. Assume that d is a positive integer such that we have

$$\text{PER}(g^d(z)) \geq \frac{1}{t}|g^d(z)| \quad \text{for all } z \in X. \quad (6)$$

Define

$$e = d + 2n - 1$$

and

$$B = \max_{z \in X} |\beta(g^e(z))|.$$

Further, assume that m is a positive integer such that

$$|g^d(z)| > 2tm \quad \text{for all } z \in X \quad (7)$$

and

$$B \leq \frac{m}{n^2 + 1}. \quad (8)$$

Define

$$L_1 = \text{COMP}(hg^e, gg^e).$$

For the proof of the next lemma see [5, Lemma 9].

Lemma 11 *Let $w \in X^*$. Then*

$$w \in L_1$$

if and only if

$$\text{Pref}_2(w) \in L_1 \text{ and } F_3(w) \subseteq F_3(L_1).$$

Now we use Lemmas 2 and 11 to show that for a word $w \in X^*$ we can decide the inclusion

$$\{g^i(w) \mid i \geq 0\} \subseteq \text{COMP}(g, h)$$

by checking whether or not $g^i(w) \in \text{COMP}(g, h)$ for a certain number of initial values of i .

Lemma 12 *Let $w \in X^*$. Then*

$$\{g^i(w) \mid i \geq 0\} \subseteq \text{COMP}(g, h)$$

if and only if

$$\{g^i(w) \mid i = 0, 1, \dots, e + 2n^2 + 2n - 3\} \subseteq \text{COMP}(g, h).$$

Proof. To prove the nontrivial part assume that $g^i(w) \in \text{COMP}(g, h)$ for $i = 0, 1, \dots, e + 2n^2 + 2n - 3$. Then $g^i(w) \in L_1$ for $i = 0, 1, \dots, 2n^2 + 2n - 3$. Hence

$$\text{Pref}_2(g^i(w)) \in L_1 \text{ and } F_3(g^i(w)) \subseteq F_3(L_1) \quad (9)$$

for $i = 0, 1, \dots, 2n^2 + 2n - 3$. Now Lemma 2 implies that (9) holds for all $i \geq 0$. Therefore Lemma 11 implies that $g^i(w) \in L_1$ for all $i \geq 0$. In other words $g^{e+i}(w) \in \text{COMP}(g, h)$ for all $i \geq 0$. Hence $g^i(w) \in \text{COMP}(g, h)$ for all $i \geq 0$. $\square$

5.2 The equality problem

In this subsection we solve the equality problem for pure morphic words generated by loop-free primitive morphisms under certain additional assumptions.

Lemma 13 *Let X be an alphabet having $n \geq 2$ letters. Define $A(n) = \lfloor 9n^3 \sqrt{n \log n} \rfloor$. Let $g : X^* \rightarrow X^*$ and $h : X^* \rightarrow X^*$ be primitive morphisms. Let $x \in X$ be a letter such that $g^\omega(x)$ and $h^\omega(x)$ exist. Assume that g is loop-free, $\text{BAL}(g, h) < \infty$ and $M_h \leq M_g^2$. Then*

$$g^\omega(x) = h^\omega(x) \quad (10)$$

if and only if

$$g^{A(n)}(x) \in \text{COMP}(g, h). \quad (11)$$

Proof. First, if (10) holds, Lemma 7 implies (11).

Assume then that (11) holds. Define the integers t, m, d, e and B as follows. First, define

$$t = 2M_g^n$$

and

$$m = \lceil (n^2 + 1)M_g^{2(2n-1)} \exp(n^2(1 + \sqrt{6n \log n})) \rceil.$$

Let q be a real number such that

$$2tm = M_g^q.$$

Then define

$$d = (\lfloor q \rfloor + 1)n, \quad e = d + 2n - 1.$$

Finally, define

$$B = \max_{z \in X} |\beta(g^e(z))|.$$

Then

$$e + 2n^2 + 2n - 2 < 9n^3 \sqrt{n \log n}. \quad (12)$$

Now we are in a position to apply Lemma 12. First, g and h are growing because g and h are primitive and $\text{card}(X) \geq 2$. Because g is loop-free, Lemma 3 implies (6). By Lemma 1 we have

$$|g^d(z)| \geq M_g^{\lfloor q \rfloor + 1} > 2tm$$

for all $z \in X$. Hence (7) holds. By Lemma 8 also (8) holds. Observe that the assumption $M_h \leq M_g^2$ implies that $M = \max\{M_g, M_h\} \leq M_g^2$. Consequently the assumptions needed for Lemma 12 hold. By (12) Lemma 12 implies that $g^i(x) \in \text{COMP}(g, h)$ for all $i \geq 0$. Now Lemma 7 implies (10). $\square$

6 The equality problem for looping primitive morphisms

In this section we solve the equality problem for pure morphic words generated by looping primitive morphisms under some additional assumptions.

Lemma 14 *Let X be an alphabet having $n \geq 2$ letters. Let $g : X^* \rightarrow X^*$ and $h : X^* \rightarrow X^*$ be primitive morphisms. Let $x \in X$ be a letter such that $g^\omega(x)$ and $h^\omega(x)$ exist. Assume that g is looping and $\text{BAL}(g, h) < \infty$. Then*

$$g^\omega(x) = h^\omega(x) \quad (13)$$

if and only if

$$g^{2n}(x) \in \text{COMP}(g, h). \quad (14)$$

Proof. Because g is looping there exist a positive integer k , a letter $z \in X$ and a nonempty primitive word v such that

$$(g^k)^\omega(z) = v^\omega.$$

Because g is primitive, each letter of X occurs in v . Because $g^k(v) \in v^*$ it follows that $g^{ki}(x)$ is a factor of v^ω for all $i \geq 0$. This implies that there is a conjugate u of v such that for infinitely many values of i , the word $g^{ki}(x)$ is a prefix of u^ω . Because $g^{ki}(x)$ is a prefix of $g^\omega(x)$ for all $i \geq 0$, we have

$$g^\omega(x) = u^\omega. \quad (15)$$

By Lemma 4 we have

$$|g^{2n}(x)| \geq 2|u|. \quad (16)$$

Because $\text{BAL}(g, h) < \infty$, we have $|g(u)| = |h(u)|$.

Assume now that (14) holds. By (15) and (16), the word u is a prefix of $g^{2n}(x)$. Hence $u \in \text{COMP}(g, h)$, which implies that $g(u) = h(u)$. Therefore $g^i(x) \in \text{COMP}(g, h)$ for all $i \geq 0$. This implies (13) by Lemma 7.

Conversely, if (13) holds, Lemma 7 implies (14). $\square$

7 The equality problem for primitive morphisms

In the previous sections we have studied the equality $g^\omega(x) = h^\omega(x)$ with the assumption that $\text{BAL}(g, h) < \infty$. It remains to give a necessary and sufficient condition for the equality $g^\omega(x) = h^\omega(x)$ without the assumption that $\text{BAL}(g, h) < \infty$.

Theorem 15 *Let X be an alphabet having $n \geq 2$ letters. Define $A(n) = \lfloor 9n^3 \sqrt{n \log n} \rfloor$. Let $g : X^* \rightarrow X^*$ and $h : X^* \rightarrow X^*$ be primitive morphisms and let $x \in X$ be a letter such that $g^\omega(x)$ and $h^\omega(x)$ exist. Define $f_1 = g^{2^{n-2}} h^{2^{n-2}}$ and $f_2 = h^{2^{n-2}} g^{2^{n-2}}$. Then*

$$g^\omega(x) = h^\omega(x) \quad (17)$$

if and only if

$$\text{BAL}(f_1, f_2) < \infty \quad (18)$$

and

$$f_1^{A(n)}(x) \in \text{COMP}(f_1, f_2). \quad (19)$$

Proof. By Lemma 1,

$$\text{alph}(g^{2^{n-2}}(z)) = \text{alph}(h^{2^{n-2}}(z)) = X \quad (20)$$

for all $z \in X$. Hence f_1 and f_2 are primitive morphisms. In particular, f_1 and f_2 are growing morphisms and $f_1^\omega(x)$ and $f_2^\omega(x)$ exist. If $y, z \in X$, then by (20) we have

$$|f_2(y)| = |h^{2^{n-2}} g^{2^{n-2}}(y)| \leq |h^{2^{n-2}} g^{2^{n-2}} h^{2^{n-2}}(z)| \leq |f_1^2(z)|.$$

Hence

$$M_{f_2} \leq M_{f_1^2} \leq M_{f_1}^2.$$

Now suppose that (17) holds. Then Lemma 6 implies that

$$f_1^\omega(x) = f_2^\omega(x). \quad (21)$$

Now Lemma 10 implies (18) and Lemma 7 implies (19).

Assume then that (18) and (19) hold. If f_1 is loop-free (resp. looping), then Lemma 13 (resp. Lemma 14) implies that (21) holds. Then we have (17) by Lemma 6. $\square$

For a method to decide the condition (18) see [4, p. 512], where it is explained how one can compute polynomials $P_1(z), \dots, P_n(z)$ depending on f_1 and f_2 such that (18) holds if and only if for each j there is a positive integer $p \leq \exp(\sqrt{6n \log n})$ such that $P_j(z)$ divides $1 - z^p$. Hence the degrees of the polynomials depend only on the cardinality of the alphabet.

References

- [1] J.-P. Allouche and J. Shallit, Automatic Sequences, Theory, Applications, Generalizations, Cambridge University Press, Cambridge, 2003.
- [2] K. Culik II and T. Harju, The ω -sequence equivalence problem for D0L systems is decidable, JACM 31 (1984) 282–298.
- [3] J. Honkala, A bound for the ω -equivalence problem of polynomial D0L systems, Theoret. Inform. Appl. 37 (2003) 149–157.
- [4] J. Honkala, An n^2 -bound for the ultimate equivalence problem of certain D0L systems over an n -letter alphabet, J. Comput. System Sci. 71 (2005) 506–519.
- [5] J. Honkala, A new bound for the D0L sequence equivalence problem, Acta Inform. 43 (2007) 419–429.
- [6] J. Honkala, The D0L ω -equivalence problem, Intern. J. Foundat. Comput. Sci. 18 (2007) 181–194.
- [7] J. Honkala, The sequence equivalence problem for primitive D0L systems, submitted.
- [8] M. Lothaire, Algebraic Combinatorics on Words, Cambridge University Press, Cambridge, 2002.
- [9] G. Rozenberg and A. Salomaa, The Mathematical Theory of L Systems, Academic Press, New York, 1980.
- [10] G. Rozenberg and A. Salomaa (Eds.), Handbook of Formal Languages, Vols. 1-3, Springer, Berlin, 1997.